



Osteoporosis and osteopenia in the distal forearm predict all-cause mortality independent of grip strength: 22-year follow-up in the population-based Tromsø Study

A. V. Hauger¹ · A. Bergland¹ · K. Holvik² · A. Ståhle^{3,4} · N. Emaus⁵ · B. H. Strand^{2,6,7}

Received: 12 April 2018 / Accepted: 30 July 2018 / Published online: 9 August 2018
© International Osteoporosis Foundation and National Osteoporosis Foundation 2018

Abstract

Summary Low bone mineral density (BMD) gives an increased risk of fractures, which can lead to premature death. Can BMD of the wrist predict mortality? BMD consistent with osteopenia and osteoporosis gave a significantly increased risk of death for both men and women in a general population in Tromsø, Norway.

Introduction To investigate if bone mineral density (BMD) levels of the distal forearm, consistent with osteopenia and osteoporosis, can predict mortality and if grip strength is an effect modifier.

Methods The study population constituted 6565 participants aged 50–79 years at baseline in the Tromsø Study wave 4 conducted in 1994–1995. Forearm BMD measured by SXA was categorized as “normal,” “osteopenia,” or “osteoporosis” following WHO’s definition. Cox regression with all-cause mortality as the outcome over 22 years of follow-up was performed for men and women separately, adjusting for health-related factors, as well as BMD by grip strength interaction. A secondary analysis with a 15-year follow-up also adjusted for hip fractures and osteoporotic fractures.

Results During follow-up, 3176 of participants died (47%). Those categorized as osteoporotic had higher mortality hazard ratio (HR) compared to those with normal BMD; men HR = 1.37 (95% confidence interval (CI) 1.19, 1.58) and women HR = 1.32 (1.14, 1.53) were adjusted for age, body mass index, physical activity, smoking habits, education, health status, chronic diseases, and grip strength. Corresponding HRs for osteopenia were men HR = 1.13 (1.00, 1.27) and women HR = 1.17 (1.01, 1.35). Further adjustments for fractures did only marginally attenuate the results, and HRs were still significant. There was no grip strength by BMD interaction.

Conclusion Men and women with low distal forearm BMD values, consistent with osteoporosis or osteopenia, had an increased mortality compared to normal BMD participants. High grip strength did not modify this association, and the association remained after adjustment for a range of health-related factors.

Keywords Bone mineral density · Grip strength · Hip fracture · Mortality · Osteoporosis · Osteopenia

✉ A. V. Hauger
anvoaha@oslomet.no

¹ Department of Physiotherapy, Faculty of Health Sciences, Oslo Metropolitan University, OsloMet – storbyuniversitetet Postboks 4 St. Olavs plass, 0130 Oslo, Norway

² Norwegian Institute of Public Health, Marcus Thranes gate 6, 0473 Oslo, Norway

³ Department of Neurobiology, Care Sciences and Society, Division of Physiotherapy, Karolinska Institutet, Alfred Nobels Allé 23, 14183 Huddinge, Sweden

⁴ Function Area Occupational Therapy & Physiotherapy, Allied Health Professionals Function, Karolinska University Hospital, Stockholm, Sweden

⁵ Department of Health and Care Sciences, Faculty of Health Sciences, UiT The Arctic University of Norway, 9037 Tromsø, Norway

⁶ Norwegian National Advisory Unit on Aging and Health, Vestfold Hospital Trust, Tønsberg, Norway

⁷ Faculty of Medicine, University of Oslo, Oslo, Norway

Introduction

Osteoporosis constitutes an important public health concern with its high incidence in Western populations, and progressive prevalence in Asia [1, 2]. Osteoporosis is known to vary by gender and age [3]. It is often defined as a disease of women because the prevalence and fracture rates are much higher among females, but once an osteoporotic hip fracture has occurred, excess mortality has been found to be higher in men [3, 4]. The incidence of osteoporosis is increasing with age, occurring mainly above the age of 50 years [5].

Osteopenia is the precursor of osteoporosis. The World Health Organization (WHO) Study Group on Osteoporosis has defined osteopenia and osteoporosis as bone mineral density (BMD) of more than 1 and 2.5 standard deviations (SDs), respectively, below the mean BMD of the young, white, female adult reference population [6]. Based on data from the USA, it has been estimated that 30% or more of all postmenopausal, white women have osteoporosis [7]. The lifetime risk of any fracture of the hip, spine, proximal or distal forearm, all considered typical osteoporotic fractures, was estimated to be 46% in women and 22% in men from age 50 years onward in a Swedish population [8]. As life expectancy increases, the population burden of osteoporosis and related fragility fractures will increase [1, 9].

A systematic review and meta-analysis from 2013 [10] found an inverse relationship between BMD and all-cause mortality. The same result was found for women with type 2 diabetes [11]. An important pathway linking low BMD to mortality is via fractures, and hip fractures in particular. Furthermore, the association between BMD and mortality could be confounded by physical fitness, physical activity, body mass, smoking habits, level of education [4, 12–15] and by comorbidity such as stroke, angina, myocardial infarction, diabetes, and asthma [16–18].

Grip strength measurements have been recommended in order to identify old people with sarcopenia [19] (low muscle mass and low muscle function). Low grip strength has also been found to predict disability, impaired quality of life, falls, and mortality [20–22], while high grip strength may indicate resilience to aging [23].

Osteoporotic fractures of the proximal femur are particularly associated with excess mortality, and studies have consistently found that this association increases with age [4, 24]. For distal forearm fractures, however, excess mortality is found to be lower or non-significant [24], but a prior wrist fracture can increase the risk of any osteoporotic fracture later in life [25, 26]. Recent studies have found that osteoporosis is more easily detected in the peripheral regions (wrist) than in the central regions (spine and hip) [27] and wrist BMD has better accuracy than lumbar BMD in diagnosing osteoporosis in postmenopausal women [28]. Measuring BMD in the distal forearm might reveal

a BMD deficiency at an earlier stage and give better prerequisites for treatment and fracture prevention.

The main aim of this paper was to assess the predictive value of established definitions for osteopenia and osteoporosis in evaluating risk of mortality. Identifying individuals at high risk is crucial in order to provide interventions on amendable risk factors for osteopenia or osteoporosis. There have been previous studies on how mortality is affected by different treatments of osteoporosis, fracture types [4, 24, 29], and BMD values in various populations [10, 11, 30]. However, the association between osteoporosis and osteopenia of the distal forearm and mortality, and the possible mediating effect of grip strength has to the very best of our knowledge not been examined in a population-based study before. Thus, an additional aim of this paper was to investigate if a strong grip modified the potential association between low BMD and mortality and whether the association was confounded by BMI, smoking, physical activity level, self-reported health status, level of education, or chronic diseases such as angina, stroke, myocardial infarction, diabetes, and asthma. We hypothesized that those with distal forearm BMD categorized as osteoporotic or osteopenic had a higher mortality risk compared to those with normal BMD values, but that this increased risk could be partly counteracted by a high grip strength.

Method

Study population

The Tromsø Study was initiated in 1974 and is a longitudinal, population-based, multi-purpose study focusing on lifestyle-related diseases [31]. There have been seven study waves, and our study population is comprised of participants from the fourth wave, conducted in 1994–1995. This wave included a bone densitometry measurement as a part of additional testing that was offered to all participants aged 55–74 years, all women aged 50–55 years, and a random selection of 10–15% of participants aged 24–55 years and 74–85 years. In the current analyses, only participants aged 50–79 years were included. The attendance rate was 76% among men and 79% among women in this age group. Our study population consisted of 6565 participants, 3818 women with a mean age of 60.7 years (SD = 7.4) and 2747 men with a mean age of 62.6 years (SD = 6.4).

Assessment of bone mineral density

Bone densitometry using SXA was performed on the non-dominant forearm at distal and ultra-distal sites with two single x-ray absorptiometry devices (DTX-100; Osteometer MediTech, Inc., Hawthorne, CA). Further specification of

the testing procedure can be found elsewhere [32]. No significant difference has been detected regarding precision of the distal and ultra-distal measurement [33]. The distal measurement was chosen for our analyses, including both radius and ulna. Osteopenia and osteoporosis were defined respectively as 1 and 2.5 SDs below the mean of young, healthy men and women (see below).

Reference values

Gender-specific internal BMD reference values were created for osteopenia and osteoporosis, based on BMD values corresponding to 1 and 2.5 standard deviations below the mean BMD of healthy men and women aged 24–39 years in the Tromsø 4 densitometry data. Besides gender and age range, the reference populations were defined by a dichotomous variable, “healthy” (yes/no), which was based on the following disease-related questions: Do you have, or have you had a myocardial infarction? (yes/no); do you have, or have you had angina pectoris? (yes/no); do you have, or have you had a cerebral stroke/brain hemorrhage? (yes/no); do you have, or have you had asthma? (yes/no); do you have, or have you had diabetes? (yes/no); and what is your current state of health? (Poor/not so good/good/very good). Those who reported “good” or “very good” self-rated health combined with “no” on all the disease-related questions were defined as “healthy,” and this group was used when calculating reference values for categorization into “normal BMD,” “osteopenia,” and “osteoporosis.” Only including the “healthy” participants resulted in 252 women with a mean BMD value of 0.471 g/cm² (SD = 0.043) and 147 men with a mean BMD value of 0.575 g/cm² (SD = 0.045). Thus, 2.5 SD below mean BMD (osteoporosis) corresponded to 0.364 g/cm² in women and 0.464 g/cm² in men, and 1.0 SD below mean (osteopenia) corresponded to 0.428 g/cm² in women and 0.531 g/cm² in men.

Ascertainment of deaths

The outcome in this study was all-cause mortality. Data on each participant was linked, by the means of the unique personal identification number, to the Norwegian Cause of Death Registry for assessment of death, and to the National Registry for assessment of emigration. Participants were followed from baseline survey in 1994–1995 until emigration, death, or November 5, 2016, whichever occurred first.

Covariates

Covariates known to be associated with lower BMD and mortality were selected a priori for inclusion as possible confounders in addition to age and gender. Height and weight was measured by trained personnel in the Tromsø Study and body mass index (BMI) was calculated as weight in kilograms

divided by height in meters squared (kg/m²) and grouped as follows: low = BMI ≤ 20.00 kg/m², normal = 20.01 kg/m² to 25.00, overweight = 25.01 to 30.00 kg/m², and obese > 30.00 kg/m². Smoking was self-reported and categorized in three groups as current, previous, or never-smoker. Education level was based on years of completed education grouped into five levels ranging from “7–10 years primary/secondary school” to “college/university 4 or more years.” Level of physical activity was self-reported by counting hours of light physical activity (not sweating or out of breath) and hard physical activity (sweating and/or out of breath) during a typical week in the previous year. The number of hours per week for each variable was categorized in four groups, i.e., none, less than one, one to two, and three or more. Chronic diseases were self-reported in Tromsø 4 with alternatives “yes” or “no” following questions about stroke, myocardial infarction, angina, diabetes, or asthma in their medical history along with questions regarding self-perceived health categorized as follows: very good, good, not so good, and poor. Grip strength of the non-dominant hand was measured in bar using a Martin vigorimeter. Each participant was allowed two attempts, and the highest score was recorded and used in analyses. Grip strength was grouped into gender-specific quartiles. Records for fractures were available for all participants until February 22, 2010. Fractures of the femur neck and trochanter were defined as “hip fractures.” These in addition to distal fractures of ulna and radius were defined as “osteoporotic fractures.” Vertebral fractures were not reported in the material.

Statistics

Separate analyses were conducted for men and women. A Cox proportional hazards survival model was used to assess the associations between T-score groups based on distal forearm BMD and mortality. We successively adjusted for health- and lifestyle-related variables in three models, i.e., model 1: (attained) age; model 2: model 1 + BMI, level of physical activity, smoking habits, and category of completed education; model 3: model 2 + self-reported health status and self-reports of chronic diseases including asthma, diabetes, angina pectoris, stroke, and myocardial infarction. In addition, grip strength by BMD interaction was tested in a fourth model. Fractures were included in a secondary analysis since fracture data was only available until February 22, 2010, giving a shorter follow-up period. Model 1 is minimally adjusted for age (attained), without fracture variables. Models 2 and 4 minimally adjusted for age (attained) and hip fractures or osteoporotic fractures. Models 3 and 5 fully adjusted in addition to hip fractures or osteoporotic fractures. The fracture variables were modeled as time-dependent covariates in order to avoid immortal time-bias. The proportional hazard (PH) assumption was inspected visually and by formal tests based on scaled Schoenfeld residuals. Statistical significance was

determined by an alpha level of 0.05. The statistical analysis was carried out with Stata/SE 15.

Results

During follow-up in the main analysis, 3176 (46.8%) of the 6790 participants died, 1538 women and 1638 men. Fifty-four participants were censored due to emigration. The mean BMD value of the total study population 50 to 79 years was 0.458 g/cm² (SD = 0.094), 0.403 g/cm² (SD 0.069) in women and 0.533 g/cm² (SD 0.067) in men (Table 1). According to the definition, 1512 (38%) female participants had normal BMD, 1329 (34%) had osteopenia, and 1104 (28%) had osteoporosis. Corresponding numbers in men were 1575 (55%), 870 (31%), and 400 (14%) (Tables 1 and 2).

In our secondary analysis including fracture data, 1242 women and 434 men experienced a fracture during a 15-year follow-up from baseline to February 22, 2010. Among women, 265 experienced a hip fracture and 479 experienced a distal forearm fracture. Corresponding numbers among men were 132 and 194.

Participants categorized as having osteoporosis were significantly older, had a lower BMI, lower grip strength, performed less hard physical activity, and had inferior self-reported health, and a higher percentage had experienced a stroke compared to those with normal BMD values (Table 2). Among women, the osteoporosis group also performed less light physical activity, they were lower educated, and had a higher lifetime prevalence of angina pectoris or a myocardial infarction than participants with normal BMD.

Significantly, more men were smokers in the osteoporosis group than in the normal BMD group.

Cox regression revealed a significantly higher mortality in both women and men with osteoporosis and osteopenia compared to the normal BMD groups (Table 3). In the fully adjusted model, including adjustments for age, BMI, level of education, physical activity, smoking, self-reported health, chronic diseases, and grip strength, the hazard ratio (HR) was 1.32 (95% confidence interval (CI) 1.14 to 1.53) for women and 1.37 (95% CI 1.19 to 1.58) for men with osteoporosis compared to those with normal BMD. Corresponding HRs for mortality in participants with osteopenia were 1.17 (95% CI 1.01 to 1.35) in women and 1.13 (95% CI 1.00 to 1.27) in men. There was no grip strength by BMD interaction in women ($p = 0.84$) or in men ($p = 0.55$), see Figs. 1 and 2 illustrating the effect of “low” (lowest quartile) and “high” (three highest quartiles) grip strength on the association between BMD as a continuous variable and HR for mortality. Tests of the proportional hazard assumption using scaled Schoenfeld residuals indicated some violation of proportionality of hazard. For osteoporosis, the HRs were comparable in the three time periods 1994–2000, 2001–2006, and 2007–2016 in both genders. For osteopenia, however, the HRs were slightly lower in the first time periods in men, while in women, they were comparable. Despite this slight violation of PH, results are presented as an average for the whole period.

In the secondary analysis, adjusting for hip fractures or osteoporotic fractures did not explain the increased mortality among participants with osteoporosis. The association between osteopenia and mortality was still significant in women after adjusting for fractures, but not in men (Table 4).

Table 1 BMD values of women and men aged 50–79 years in the Tromsø 4 Study, categorized as “normal,” “osteopenia,” and “osteoporosis” and number of deaths within age groups of 5 years

Age group	<i>N</i>	Mean BMD (SD)	% normal	% osteopenia	% osteoporosis	<i>n</i> (%) deaths
Women	3945	0.403 (0.069)	38.3	33.7	28.0	1538 (39.0)
50–54	1050	0.453 (0.050)	70.2	25.3	4.5	135 (12.9)
55–59	840	0.421 (0.055)	46.1	39.5	14.4	175 (20.8)
60–64	695	0.393 (0.060)	27.5	41.9	30.7	247 (35.5)
65–69	752	0.365 (0.064)	16.4	34.7	48.9	475 (63.2)
70–74	577	0.352 (0.063)	12.0	29.8	58.2	477 (82.7)
75–79	31	0.341 (0.081)	16.1	22.6	61.3	29 (93.5)
Men	2845	0.533 (0.067)	55.4	30.6	14.0	1638 (57.6)
50–54	225	0.564 (0.050)	77.8	20.0	2.2	40 (17.8)
55–59	793	0.552 (0.056)	67.6	27.1	5.3	258 (32.5)
60–64	700	0.539 (0.061)	56.1	32.7	11.1	378 (54.0)
65–69	606	0.520 (0.069)	46.4	33.7	20.0	468 (77.2)
70–74	494	0.501 (0.073)	36.0	34.4	29.6	467 (94.5)
75–79	27	0.492 (0.099)	44.4	25.9	29.6	27 (100.0)

N total number of participants, *BMD* bone mineral density, *n* number of participants who died during the follow-up period

Table 2 Number or proportions (%) of participants in the Tromsø 4 Study in groups: “normal BMD,” “osteopenia,” and “osteoporosis” at baseline in 1994/1995. Number of deaths, person years (py), and mortality rate per 1000 py during a 22-year follow-up. Each variable listed in women and men

	<i>N</i>	Normal BMD	Osteopenia	Osteoporosis	Trend*
Women					
Number of participants	3945	1512	1329	1104	
Number of deaths	1501	327	498	676	< 0.001
Person years (py)	3945	30,032	24,572	18,112	
Mortality rate per 1000 py	3945	10.9	20.3	37.3	
Age (years)	3945	56.5 (6.1)	61.1 (6.8)	66.1 (5.8)	< 0.001
BMD (g/cm ²)	3945	0.472 (0.032)	0.398 (0.019)	0.316 (0.037)	< 0.001
BMI (kg/m ²)	3937	26.6 (4.5)	26.2 (4.5)	25.2 (4.4)	< 0.001
Grip strength (bar)	3931	0.79 (0.20)	0.73 (0.18)	0.66 (0.18)	< 0.001
Smoking	3941				0.138
Never smoker (%)		43.2	45.4	40.9	–
Current smoker (%)		30.2	27.9	32.5	–
Previous smoker (%)		26.6	26.7	26.7	–
L-phys.act. < 1 h/week (%)	3938	25.4	26.3	29.6	0.044
H-phys.act. < 1 h/week (%)	3903	81.9	86.0	90.4	< 0.001
Low education (%)	3915	55.2	60.8	70.2	< 0.001
Self-reported health status	3939				< 0.001
Poor (%)		2.5	3.1	4.9	–
Not so good (%)		42.6	46.5	52.5	–
Good (%)		47.9	44.6	38.9	–
Very good (%)		7.0	5.9	3.6	–
Stroke (%)	3929	1.2	2.1	2.5	0.042
Angina (%)	3936	4.6	6.4	10.1	< 0.001
Myocardial infarction (%)	3932	1.9	3.2	4.0	0.004
Diabetes (%)	3929	2.3	3.5	3.2	0.132
Asthma (%)	3927	8.0	8.9	9.0	0.577
Men					
Number of participants	2845	1575	870	400	
Number of deaths	1596	740	530	326	< 0.001
Person years (py)	2845	26,944	13,760	5068	
Mortality rate per 1000 py	2845	27.5	38.5	64.3	
Age (years)	2845	61.2 (6.2)	63.4 (6.1)	66.8 (5.6)	< 0.001
BMD (g/cm ²)	2845	0.581 (0.036)	0.501 (0.019)	0.419 (0.039)	< 0.001
BMI (kg/m ²)	2843	26.6 (3.2)	25.6 (3.3)	24.9 (3.8)	< 0.001
Grip strength (bar)	2831	0.89 (0.21)	0.83 (0.20)	0.72 (0.19)	< 0.001
Smoking					< 0.001
Never smoker (%)	2844	20.9	17.5	12.8	–
Current smoker (%)	2844	27.4	33.0	40.8	–
Previous smoker (%)	2844	51.7	49.5	46.5	–
L-phys.act. < 1 h/week (%)	2828	25.1	25.0	27.5	0.603
H-phys.act. < 1 h/week (%)	2814	70.9	74.3	78.4	0.006
Low education (%)	2833	47.2	45.9	52.3	0.131
Self-reported health status	2841				0.014
Poor (%)		3.2	2.8	6.0	–
Not so good (%)		39.8	38.4	42.7	–
Good (%)		50.9	53.4	47.7	–
Very good (%)		6.1	5.4	3.5	–
Stroke (%)	2834	2.7	3.2	6.3	0.002

Table 2 (continued)

	<i>N</i>	Normal BMD	Osteopenia	Osteoporosis	Trend*
Angina (%)	2836	12.2	12.2	14.3	0.515
Myocardial infarction (%)	2836	9.8	10.7	12.3	0.325
Diabetes (%)	2834	3.8	3.6	3.0	0.779
Asthma (%)	2833	7.4	6.1	9.1	0.161

"< 1 h/week" contains both alternatives "none" and "less than one." "Low education" = 7 years or less

L/H-phys.act. light/hard physical activity, *BMD* bone mineral density, *py* person years

*Trend gives *p* values based on linear regression for the continuous variables (normal BMD coded 0, osteopenia coded 1, and osteoporosis coded 2) and chi-square test for the categorical ones

Discussion

To the best of our knowledge, this is the first population-based study to examine the association between osteoporosis and osteopenia of the distal forearm and mortality and the possible mediating effect of grip strength. We found a statistically significant association between osteopenic and osteoporotic BMD levels of the distal forearm and increased mortality rate in both women and men.

The strengths of the present study include the population-based design, standardized objective measures of bone mineral density and grip strength, a large sample size, and a long follow-up of 22 years with updated time of death from as recently as November 2016. The population consists of people

living in both rural and urban areas, and the study had a high attendance rate (about 78%).


However, the study is not without limitations. Self-reported variables challenge the internal validity of any study [34, 35]. State of health, presence of chronic diseases, level of physical activity, education, and smoking habits are self-reported variables and might be subject to over- or under-estimation due to recall bias [36] or socially desirable responding (SDR) [37]. This can in turn lead to an under-estimation of the potential association between variables. Though this could be the case with some of the variables mentioned above, the outcome in the current analysis was the registry-based hard endpoint of deaths, while our main exposure variables (BMD and grip strength) were measured objectively.

Table 3 Hazard ratios (HR) with 95% confidence intervals (CI) of mortality for BMD categories: "normal," "osteopenia," and "osteoporosis" during a 22-year follow-up from 1994 to 1995 to November 2016. Models 1–4 progressively adjusted for age and lifestyle- and health-related covariates

Model adjusted for	BMD category	<i>N</i>	Women			Men		
			HR	95% CI		HR	95% CI	
Model 1: age	Normal	3818	1.00			1.00		
	Osteopenia		1.17	1.01	1.35	1.14	1.02	1.28
	Osteoporosis		1.42	1.23	1.64	1.62	1.41	1.85
						2747		
Model 2: model 1 + BMI, education, physical activity, smoking	Normal	3818	1.00			1.00		
	Osteopenia		1.17	1.01	1.36	1.13	1.01	1.27
	Osteoporosis		1.35	1.17	1.57	1.45	1.26	1.67
						2747		
Model 3: model 2 + self-reported health and chronic diseases*	Normal	3818	1.00			1.00		
	Osteopenia		1.18	1.02	1.36	1.14	1.01	1.28
	Osteoporosis		1.34	1.16	1.55	1.42	1.23	1.64
						2747		
Model 4: model 3 + grip strength	Normal	3818	1.00			1.00		
	Osteopenia		1.17	1.01	1.35	1.13	1.00	1.27
	Osteoporosis		1.32	1.14	1.53	1.37	1.19	1.58
						2747		

*Chronic diseases include angina, asthma, stroke, myocardial infarction, and diabetes. *N* number of subjects included in analysis

Fig. 1 Mortality hazard ratios with 95% confidence intervals across the range of distal forearm BMD in women with low grip strength (dashed curve) and in women with normal or high grip strength (solid curve). Mediating effect not significant



We controlled for variables that were measured at baseline in 1994/5. During the follow-up of 22 years, it is likely that some variables changed, especially the presence of chronic diseases since it is well-known that comorbidity increases in older age. The participants may also have experienced significant changes in BMD during follow-up that could be associated with excess mortality. This could be subject for further research.

We created our own reference values in order to define osteopenia and osteoporosis for our population, but the association between BMD as a continuous variable and mortality was also analyzed, allowing the reader to study the whole spectrum of BMD independent of our categorization into

osteoporosis, osteopenia, and normal BMD. Modern methods for BMD testing have changed over the past 22 years, and we were unsuccessful in retrieving external reference values for SXA of the distal forearm. There are both strengths and limitations in creating our own reference values. We have no guarantee that our reference groups are similar to those used in other studies, and the variation within the reference group warrants the size of 1 SD which in turn make out the cutoff values. However, this resulted in 28% of the women being categorized as osteoporotic and this is comparable to other findings in Caucasian women [7], considering that the oldest old were not included in this study. A strength of creating a reference group from the same study is that they share the

Fig. 2 Mortality hazard ratios with 95% confidence intervals across the range of distal forearm BMD in men with low grip strength (dashed curve) and in women with normal or high grip strength (solid curve). Mediating effect not significant




Table 4 Hazard ratios (HR) with 95% confidence intervals (CI) of mortality for BMD categories: “normal,” “osteopenia,” and “osteoporosis” and after sustaining a hip fracture or osteoporotic fracture during a 15-year follow-up from 1994 to 1995 to March 2010. Model 1 minimally adjusted without fractures. Models 2 and 4 minimally adjusted and models 3 and 5 fully adjusted for age and lifestyle- and health-related covariates in addition to hip fractures (model 2 and 3) and osteoporotic fractures (model 4 and 5)

Model adjusted for	BMD category	N	Women			N	Men		
			HR	95% CI			HR	95% CI	
Model 1: age	Normal	3809	1.00			2745	1.00		
	Osteopenia		1.25	1.02	1.54		1.06	0.92	1.23
	Osteoporosis		1.46	1.19	1.79		1.54	1.30	1.82
Model 2: model 1 + hip fracture	Normal	3809	1.00			2745	1.00		
	Osteopenia		1.24	1.01	1.53		1.05	0.90	1.21
	Osteoporosis		1.40	1.14	1.72		1.52	1.28	1.79
Model 3: model 2 + BMI, education, physical activity, smoking, self-reported health, grip strength, and chronic diseases*	Normal	3809	1.00			2745	1.00		
	Osteopenia		1.24	1.01	1.52		1.00	0.86	1.16
	Osteoporosis		1.30	1.05	1.61		1.25	1.05	1.49
Model 4: age, osteoporotic fracture	Normal	3809	1.00			2745	1.00		
	Osteopenia		1.24	1.01	1.52		1.04	0.90	1.21
	Osteoporosis		1.42	1.15	1.74		1.50	1.27	1.77
Model 5: model 4 + BMI, education, physical activity, smoking, self-reported health, grip strength, and chronic diseases*	Normal	3809	1.00			2745	1.00		
	Osteopenia		1.23	1.00	1.51		1.00	0.86	1.17
	Osteoporosis		1.30	1.05	1.61		1.23	1.04	1.47

*Chronic diseases include angina, asthma, stroke, myocardial infarction and diabetes. *N* number of subjects included in analysis

same geographical and cultural affiliation; we know how the BMD has been measured and tests are performed by the same professionals, following the same protocols as in the main analyses.

Dementia and other cognitive impairments increase the risk of mortality. 6.1% of all deaths in Norway in 2016 were registered with dementia as the underlying cause of death [38]. Cognitive assessments were not incorporated in Tromsø 4 so we could not control for cognitive impairments or dementia at baseline in our analysis; however, later study waves including the same population revealed that a low proportion of the participants had cognitive impairments, with 7.3% scoring low on one or more of the cognitive tests in addition to self-report of memory problems. Out of these, only one participant had dementia. It is therefore unlikely that dementia confounded the association we found between osteoporosis/osteopenia and mortality.

Our findings indicate that BMD measurements of the distal forearm has a predictive value in mortality risk assessment, and mortality can be predicted using the commonly accepted T-values of -1 and -2.5 for osteopenia and osteoporosis, though only demonstrated on a group level. In evaluating an

individual's mortality risk, osteopenia and osteoporosis should be viewed as independent risk factors of death that will add to the total risk along with other known risk factors.

The association between mortality and osteoporosis was slightly stronger in men while the association with osteopenia was somewhat stronger in women, indicating that smaller deficiencies in BMD might be more serious in women. However, the between-gender differences are not large enough to make such assumptions based on this material. The association between osteoporosis and increased mortality was still significant in both men and women after adjusting for fractures, indicating that there might be a more complex relationship between low BMD and mortality risk than we are currently aware of. Several authors have found an inverse relationship between BMD and risk of cardiovascular disease and cardiovascular death [39, 40]. Although we controlled for myocardial infarction and angina, these variables were measured at baseline and more cases probably occurred during follow-up, potentially more often among those with low BMD.

That our main analysis also shows significantly higher mortality for osteopenic BMD values suggests that it might

be valuable to initiate treatment measures already at this stage, though previous research debates the cost-effectiveness of pharmacological treatment of osteopenia purely based on T-scores [41, 42]. Low BMD is mainly seen as a risk factor of fractures, and it has been debated whether expensive medication is the right way to prevent fractures as opposed to means of fall prevention [43]. However, one intervention does not exclude the other, and fall prevention should be emphasized regardless of any medical prescriptions. In Norway, osteoporosis appears to be both under-diagnosed and under-treated according to Devold et al. [44] who found that 1 year after experiencing a hip fracture, only 14.6% of women and 4.2% of men used some form of anti-osteoporotic medication. Gray et al. [29] found a significant reduction of mortality risk associated with use of fracture-preventing medication in their meta-analysis, and the effect was largest in older, frailer individuals. The decision to prescribe medication should in any case be based on a full assessment of the person's fracture risk and potential benefits of treatment.

In our study, a general population was screened for low BMD independent of prior indication of a BMD deficiency. There are currently no routines for general screening of BMD in Norway, but our findings indicate that general BMD measurements can be of value in identifying individuals with higher risk of mortality. Schousboe et al. [45] found that universal BMD screening of the population combined with alendronate therapy for those found to have osteoporosis is highly cost-effective for women aged 65 and older and may be cost-saving for ambulatory women aged 85 and older.

Based on our study, we cannot conclude whether treatment of low BMD will help decrease risk of mortality or if the BMD deficiency is merely a marker for frailty. In practical terms, measured osteopenia and osteoporosis in the distal forearm reveals individuals with increased risk of mortality, which warrants closer follow-up of these individuals by health care personnel.

In a previous analysis from the Tromsø 4 Study wave, high grip strength was associated with lower risk of mortality [20], yet grip strength did not attenuate or modify the higher mortality risk for participants with osteoporosis or osteopenia in our analyses. Thus, these variables seem to be independently associated with mortality.

In elderly people, most wrist fractures occur in individuals with low BMD who are relatively healthy and active and have good neuromuscular function [46]. BMD is commonly measured after a low-energy trauma fracture. Even though a wrist fracture in itself has not been found to increase the risk of mortality [24], our findings indicate that an underlying BMD deficiency in the forearm can have more serious implications, and measures should be taken accordingly with respect to current medical guidelines for prevention of fractures and treatment of osteoporosis.

Conclusion

Women and men with distal forearm BMD values consistent with both osteoporosis and osteopenia had an increased all-cause mortality compared to people with normal BMD values, independent of lifestyle- and health-related variables. The association between BMD and all-cause mortality was not modified by hand grip strength.

Funding No external funding has been received to finance this project. The manuscript will be part of Annette V. Hauger's Ph.D thesis and is internally funded by Oslo Metropolitan University.

Compliance with ethical standards

Conflict of interest None.

References

- Cooper C, Cole Z, Holroyd C, Earl S, Harvey N, Dennison E, Melton L, Cummings S, Kanis J (2011) Secular trends in the incidence of hip and other osteoporotic fractures. *Osteoporos Int* 22: 1277–1288
- Cauley JA (2013) Public health impact of osteoporosis. *J Gerontol A Biomed Sci Med Sci* 68:1243–1251
- Pietschmann P, Rauner M, Sipos W, Kersch-Schindl K (2009) Osteoporosis: an age-related and gender-specific disease—a mini-review. *Gerontology* 55:3–12
- Omsland TK, Emaus N, Tell GS, Magnus JH, Ahmed LA, Holvik K, Center J, Forsmo S, Gjesdal CG, Schei B (2014) Mortality following the first hip fracture in Norwegian women and men (1999–2008). A NOREPOS study. *Bone* 63:81–86
- Bor A, Matuz M, Gyimesi N, Biczók Z, Soós G, Doró P (2015) Gender inequalities in the treatment of osteoporosis. *Maturitas* 80: 162–169
- Kanis JA (1994) Assessment of fracture risk and its application to screening for postmenopausal osteoporosis: synopsis of a WHO report. *Osteoporos Int* 4:368–381
- Kanis JA, Melton LJ, Christiansen C, Johnston CC, Khaltaev N (1994) The diagnosis of osteoporosis. *J Bone Miner Res* 9:1137–1141
- Kanis J, Johnell O, Oden A, Sernbo I, Redlund-Johnell I, Dawson A, De Laet C, Jonsson B (2000) Long-term risk of osteoporotic fracture in Malmö. *Osteoporos Int* 11:669–674
- Omsland T, Magnus J (2014) Forecasting the burden of future postmenopausal hip fractures. *Osteoporos Int* 25:2493–2496
- Qu X, Huang X, Jin F, Wang H, Hao Y, Tang T, Dai K (2013) Bone mineral density and all-cause, cardiovascular and stroke mortality: a meta-analysis of prospective cohort studies. *Int J Cardiol* 166:385–393
- Lenchik L, Register TC, Hsu F-C, Xu J, Smith SC, Carr JJ, Freedman BI, Bowden DW (2017) Bone mineral density of the radius predicts all-cause mortality in patients with type 2 diabetes: diabetes heart study. *J Clin Densitom*
- Fry PS, Debats DL (2006) Sources of life strengths as predictors of late-life mortality and survivorship. *Int J Aging Hum Dev* 62:303–334
- Frith PA (2005) Impact of smoking, diabetes and hypertension on survival in the elderly: the Dubbo study. *Med J Aust* 182:495

14. Brown DC, Hayward MD, Montez JK, Hummer RA, Chiu C-T, Hidajat MM (2012) The significance of education for mortality compression in the United States. *Demography* 49:819–840
15. Wong PK, Christie JJ, Wark JD (2007) The effects of smoking on bone health. *Clin Sci* 113:233–241
16. Loprinzi PD, Addoh O (2017) Accelerometer-determined physical activity and all-cause mortality in a National Prospective Cohort Study of Adults Post-Acute Stroke. *Am J Health Promot* 0890117117720061
17. Zahra A, Lee E-W, L-y S, Park J-H (2015) Cardiovascular disease and diabetes mortality, and their relation to socio-economical, environmental, and health behavioural factors in worldwide view. *Public Health* 129:385–395
18. Ebmeier S, Thayabaran D, Braithwaite I, Bénamara C, Weatherall M, Beasley R (2017) Trends in international asthma mortality: analysis of data from the WHO Mortality Database from 46 countries (1993–2012). *Lancet* 390:935–945
19. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, Martin FC, Michel J-P, Rolland Y, Schneider SM (2010) Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing* 39:412–423
20. Strand BH, Cooper R, Bergland A, Jørgensen L, Schirmer H, Skirbekk V, Emaus N (2016) The association of grip strength from midlife onwards with all-cause and cause-specific mortality over 17 years of follow-up in the Tromsø Study. *J Epidemiol Community Health* 70:1214–1221
21. Sayer AA, Syddall HE, Martin HJ, Dennison EM, Anderson FH, Cooper C (2006) Falls, sarcopenia, and growth in early life: findings from the Hertfordshire cohort study. *Am J Epidemiol* 164:665–671
22. Syddall HE, Martin HJ, Harwood RH, Cooper C, Sayer AA (2009) The SF-36: a simple, effective measure of mobility-disability for epidemiological studies. *JNHA J Nutr Health Aging* 13:57–62
23. Rantanen T, Masaki K, He Q, Ross GW, Willcox BJ, White L (2012) Midlife muscle strength and human longevity up to age 100 years: a 44-year prospective study among a decedent cohort. *Age* 34:563–570
24. Leboime A, Confavreux CB, Mehse N, Paccou J, David C, Roux C (2010) Osteoporosis and mortality. *Joint Bone Spine* 77:S107–S112
25. Barrett-Connor E, Sajjan S, Siris E, Miller P, Chen Y-T, Markson L (2008) Wrist fracture as a predictor of future fractures in younger versus older postmenopausal women: results from the National Osteoporosis Risk Assessment (NORA). *Osteoporos Int* 19:607–613
26. Ahmed LA, Center JR, Bjørnerem Å, Bluic D, Joakimsen RM, Jørgensen L, Meyer HE, Nguyen ND, Nguyen TV, Omsland TK (2013) Progressively increasing fracture risk with advancing age after initial incident fragility fracture: the Tromsø study. *J Bone Miner Res* 28:2214–2221
27. Abdelmohsen AM (2017) Comparison of central and peripheral bone mineral density measurements in postmenopausal women. *J Chiropr Med* 16:199–203
28. Eftekhari-Sadat B, Ghavami M, Toopchizadeh V, Ghahvechi Akbari M (2016) Wrist bone mineral density utility in diagnosing hip osteoporosis in postmenopausal women. *Ther Adv Endocrinol Metab* 7:207–211
29. Grey A, Bolland M (2013) The effect of treatments for osteoporosis on mortality. *Osteoporosis International* 1–6
30. Miyake H, Kanazawa I, Sugimoto T (2018) Association of bone mineral density, bone turnover markers, and vertebral fractures with all-cause mortality in type 2 diabetes mellitus. *Calcif Tissue Int* 102: 1–13
31. Jacobsen BK, Eggen AE, Mathiesen EB, Wilsgaard T, Njølstad I (2011) Cohort profile: the Tromsø Study. *Int J Epidemiol* 41:961–967
32. Berntsen GKR, Fønnebo V, Tollan A, Sjøgaard AJ, Magnus JH (2001) Forearm bone mineral density by age in 7,620 men and women the Tromsø Study, a population-based study. *Am J Epidemiol* 153
33. Berntsen GKR, Fønnebo V, Tollan A, Sjøgaard AJ, Joakimsen RM, Magnus JH (2000) The Tromsø Study: determinants of precision in bone densitometry. *J Clin Epidemiol* 53:1104–1112
34. Prince SA, Adamo KB, Hamel ME, Hardt J, Gorber SC, Tremblay M (2008) A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 5:56
35. Gorber SC, Tremblay M, Moher D, Gorber B (2007) A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev* 8:307–326
36. Hassan E (2006) Recall bias can be a threat to retrospective and prospective research designs. *Internet J Epidemiol* 3:339–412
37. Van de Mortel TF (2008) Faking it: social desirability response bias in self-report research. *Aust J Adv Nurs* 25:40
38. Norwegian Cause of Death Registry (2017) Dødsårsaksregisteret - Statistikkbank. Norwegian Institute of Public Health <http://statistikkbank.fhi.no/dar/>. Accessed 28 June 2018
39. Marcovitz PA, Tran HH, Franklin BA, O'Neill WW, Yerkey M, Boura J, Kleerekoper M, Dickinson CZ (2005) Usefulness of bone mineral density to predict significant coronary artery disease. *Am J Cardiol* 96:1059–1063
40. von der Recke P, Hansen MA, Hassager C (1999) The association between low bone mass at the menopause and cardiovascular mortality. *Am J Med* 106:273–278
41. McClung MR (2005) Osteopenia: to treat or not to treat? *Ann Intern Med* 142:796–797
42. Seeman E (2004) Treatment of osteoporosis—why, whom, when and how to treat. *Indian J Endocrinol Metab* 6:60
43. Järvinen TL, Sievänen H, Khan KM, Heinonen A, Kannus P (2008) Shifting the focus in fracture prevention from osteoporosis to falls. *Bmj* 336:124–126
44. Devold H, Sjøgaard A, Tverdal A, Falch J, Furu K, Meyer H (2013) Hip fracture and other predictors of anti-osteoporosis drug use in Norway. *Osteoporos Int* 24:1225–1233
45. Schousboe JT, Ensrud KE, Nyman JA, Melton LJ, Kane RL (2005) Universal bone densitometry screening combined with alendronate therapy for those diagnosed with osteoporosis is highly cost-effective for elderly women. *J Am Geriatr Soc* 53:1697–1704
46. Kelsey JL, Browner WS, Seeley DG, Nevitt MC, Cummings SR, Group SoOFR (1992) Risk factors for fractures of the distal forearm and proximal humerus. *Am J Epidemiol* 135:477–489